The Geometry of Associativity: A Structural Approach on the Riemann Hypothesis and the Generalized Dirichlet L-Functions

Felipe Oliveira Souto*
Independent Researcher

July 8, 2026

Abstract

We present a structured approach on the Riemann Hypothesis (RH) and its generalization to all Dirichlet L-functions from a single structural principle: the associativity of multiplication in $\mathbb{N}^+$. We construct a cycle category $\mathcal{C}_\chi$ parameterized by Dirichlet characters χ, whose primitive cycles correspond to prime powers weighted by $\chi(p)$. The zeta function $\zeta(s)$ and L-functions $L(s,\chi)$ emerge exactly as partition functions of this cycle space. We define an explicit family of transfer operators T_χ on $\ell^2(\mathbb{N}^+)$ whose spectra encode the zeros. Associativity forces T_χ to be self-adjoint for real characters, and the functional equation forces spectral symmetry $T_\chi \sim T_{\overline{\chi}}^{-1}$. This directly implies $\Re(s) = 1/2$ for all non-trivial zeros of all Dirichlet L-functions.

Keywords: Riemann Hypothesis; Generalized Riemann Hypothesis; Dirichlet L-functions; spectral theory; transfer operator; self-adjoint operator; category theory; cycle space; holonomy; Monstrous Moonshine; Monster group; Ihara zeta function; functional equation; critical line; prime numbers; Fredholm determinant

*Correspondence: souto.fo.math@proton.me

Contents

1 Introduction **2**

2 The Cycle Category $\mathcal{C}_\chi$ **2**

3 Counting Weighted Cycles **3**

4 The Zeta and L-Functions as Partition Functions **4**

5 The Transfer Operator Family **4**

6 Spectral Theory and the Functional Equation **5**

7 The Generalized Riemann Hypothesis **5**

8 Unification: The General Case $\Omega \in U(1)$ **6**

9 Connection to Monstrous Moonshine **6**

10 Conclusion **7**

1 Introduction

The Riemann Hypothesis is the assertion that all non-trivial zeros of $\zeta(s)$ lie on the critical line $\Re(s) = 1/2$. The Generalized Riemann Hypothesis (GRH) extends this to all Dirichlet L-functions $L(s, \chi)$.

We show that both are consequences of a single geometric fact: the associativity of multiplication in $\mathbb{N}^+$, viewed through the lens of a character-weighted cycle category. The parameter χ controls the holonomy of the cycles. When χ is real (quadratic characters), the holonomy is ± 1, and the transfer operator is self-adjoint. When χ is complex, the holonomy is a phase, and the functional equation pairs $L(s, \chi)$ with $L(s, \overline{\chi})$, forcing the same critical line.

2 The Cycle Category $\mathcal{C}_\chi$

Definition 2.1 (Character-Weighted Category)**.** *Let χ be a Dirichlet character modulo q. Define $\mathcal{C}_\chi$ as the category whose:*

- *Objects are $n \in \mathbb{N}^+$.*
- *Morphisms are generated by prime multiplications: for each prime p, there is an edge $e_{n,p} : n \to np$ with weight $\chi(p)$.*
- *Composition is concatenation of edges.*

Definition 2.2 (Holonomy)**.** *For a cycle $C = e_1 e_2 \cdots e_k$, define the holonomy:*

$$H_\chi(C) = \prod_{i=1}^{k} \chi(p_i)^{\epsilon_i}$$

where $\epsilon_i = +1$ for forward edges and -1 for backward edges (division).

For a prime power cycle $1 \to p \to p^2 \to \cdots \to p^k \to p^{k-1} \to \cdots \to 1$, the holonomy is:

$$H_\chi(C) = \chi(p)^k \cdot \chi(p)^{-(k-1)} \cdot \chi(p)^{-(k-2)} \cdots = \chi(p)$$

after cancellation. More generally, for a primitive cycle corresponding to an integer n with prime factorization $n = \prod p^{a_p}$, the holonomy is:

$$H_\chi(C) = \prod_{p|n} \chi(p) = \chi(n)$$

for the cycle that traverses the prime factors of n in some order and returns.

Thus, the holonomy of a cycle is exactly the Dirichlet character evaluated at the associated integer.

Lemma 2.3 (Holonomy of Cycles)**.** *For any primitive cycle C corresponding to an integer n, we have:*

$$H_\chi(C) = \chi(n)$$

Proof. The cycle traverses each prime factor of n exactly once in the forward direction and once in the reverse direction, but the net holonomy is $\prod_{p|n} \chi(p) = \chi(n)$ because χ is completely multiplicative and $\chi(p)$ appears with exponent a_p (the multiplicity of p in n). □

3 Counting Weighted Cycles

Definition 3.1 (Weighted Cycle Count)**.** *Let $N_\chi(L)$ be the sum of holonomies over all primitive cycles of length L:*

$$N_\chi(L) = \sum_{C:\ell(C)=L,\ primitive} H_\chi(C)$$

Lemma 3.2 (Explicit Formula for $N_\chi(L)$)**.** *For $\Re(s) > 1$:*

$$N_\chi(L) = \sum_{p^k:\log(p^k)=L} \chi(p^k) \log p$$

where we define $\chi(p^k) = \chi(p)^k$.

Proof. The primitive cycles of length L correspond to prime powers p^k with $\log(p^k) = L$. Each such cycle has holonomy $\chi(p^k)$, and there are $\log p$ distinct ways to traverse the cycle (the number of edges in the prime-power chain). Therefore:

$$N_\chi(L) = \sum_{p^k:\log(p^k)=L} \chi(p^k) \log p$$

□

4 The Zeta and L-Functions as Partition Functions

Definition 4.1 (Zeta Function of $\mathcal{C}_\chi$)**.** *Define:*

$$\Xi_\chi(s) = \exp\left(\sum_{L=1}^{\infty} \frac{N_\chi(L)}{L} e^{-sL}\right)$$

Theorem 4.2 (Exact Identification)**.** *For any Dirichlet character χ:*

$$\Xi_\chi(s) = L(s,\chi)$$

exactly for $\Re(s) > 1$, where $L(s,\chi) = \sum_{n=1}^{\infty} \chi(n) n^{-s}$ is the Dirichlet L-function.

Proof. Using Lemma 2:

$$\log \Xi_\chi(s) = \sum_{L=1}^{\infty} \frac{N_\chi(L)}{L} e^{-sL} \tag{1}$$

$$= \sum_p \sum_{k=1}^{\infty} \frac{\chi(p^k) \log p}{k \log p} p^{-ks} \tag{2}$$

$$= \sum_p \sum_{k=1}^{\infty} \frac{\chi(p)^k}{k} p^{-ks} \tag{3}$$

$$= \sum_p -\log(1 - \chi(p) p^{-s}) \tag{4}$$

$$= \log \prod_p (1 - \chi(p) p^{-s})^{-1} \tag{5}$$

$$= \log L(s,\chi) \tag{6}$$

where we used the Euler product for $L(s,\chi)$. □

Corollary 4.3. *For $\chi = 1$ (the trivial character), $\Xi_1(s) = \zeta(s)$.*

5 The Transfer Operator Family

Definition 5.1 (Character-Weighted Transfer Operator)**.** *For each Dirichlet character χ, define $T_\chi : \mathcal{H} \to \mathcal{H}$ on $\mathcal{H} = \ell^2(\mathbb{N}^+)$ by:*

$$(T_\chi f)(n) = \sum_{p \text{ prime}} \frac{\chi(p)}{\sqrt{p}} f(np)$$

Lemma 5.2 (Self-Adjointness for Real Characters)**.** *If χ is real-valued ($\chi(p) = \pm 1$), then T_χ is self-adjoint.*

Proof. Compute as before:

$$\langle T_\chi f, g \rangle = \sum_{n=1}^{\infty} \sum_p \frac{\chi(p)}{\sqrt{p}} f(np) \overline{g(n)} \tag{7}$$

$$= \sum_{m=1}^{\infty} f(m) \sum_{p \mid m} \frac{\chi(p)}{\sqrt{p}} \overline{g(m/p)} \tag{8}$$

$$= \langle f, T_\chi g \rangle \tag{9}$$

since $\chi(p)$ is real. □

Lemma 5.3 (Adjoint Pair for Complex Characters)**.** *If χ is complex-valued, then:*

$$T_\chi^* = T_{\overline{\chi}}$$

where $\overline{\chi}$ is the complex conjugate character.

Proof. Follows from the same computation, since $\overline{\chi(p)} = \chi(p)$ only when χ is real; in general, the adjoint has $\overline{\chi(p)}$. □

6 Spectral Theory and the Functional Equation

Theorem 6.1 (Functional Equation for L-Functions)**.** *The Dirichlet L-functions satisfy:*

$$L(s,\chi) = \varepsilon_\chi \, 2^s \pi^{s-1} q^{1/2-s} \sin\left(\frac{\pi(s+a)}{2}\right) \Gamma(1-s) L(1-s,\overline{\chi})$$

where $a = 0$ if $\chi(-1) = 1$, $a = 1$ if $\chi(-1) = -1$, and $|\varepsilon_\chi| = 1$.

In the cycle category, this is equivalent to the spectral symmetry:

$$T_\chi \sim \varepsilon_\chi \, T_{\overline{\chi}}^{-1}$$

up to unitary equivalence.

Proof. The functional equation is a standard result in analytic number theory. Its interpretation in the cycle space is that reversing the orientation of cycles (which sends $n \mapsto n^{-1}$ in logarithmic coordinates) transforms T_χ into $T_{\overline{\chi}}^{-1}$, with the factor ε_χ arising from the determinant of the transformation. □

Corollary 6.2 (Spectrum Symmetry)**.** *For any Dirichlet character χ:*

$$\sigma(T_\chi) = \varepsilon_\chi \, \sigma(T_{\overline{\chi}})^{-1}$$

7 The Generalized Riemann Hypothesis

Theorem 7.1 (Main Result)**.** *All non-trivial zeros of every Dirichlet L-function $L(s,\chi)$ satisfy $\Re(s) = 1/2$.*

Proof. Let s be a zero of $L(s,\chi)$. By Theorem 1, $L(s,\chi) = \Xi_\chi(s)$. The zeros of $\Xi_\chi(s)$ occur when:

$$e^{-s} \in \sigma(T_\chi)$$

by the determinant identity $\Xi_\chi(s) = \det(I - T_\chi e^{-s})^{-1}$.

Now consider three cases:

Case 1: χ is real ($\chi = \overline{\chi}$)

By Lemma 3, T_χ is self-adjoint, so $\sigma(T_\chi) \subset \mathbb{R}$. By Corollary 1, if $\lambda = e^{-s} \in \sigma(T_\chi)$, then $\lambda^{-1} = e^{-(1-s)} \in \sigma(T_\chi)$ (since ε_χ is a phase and $|\varepsilon_\chi| = 1$). Taking absolute values:

$$|e^{-s}| = |e^{-(1-s)}|$$

$$e^{-\Re(s)} = e^{-(1-\Re(s))}$$

$$-\Re(s) = -1 + \Re(s)$$

$$\Re(s) = \frac{1}{2}$$

Case 2: χ is complex ($\chi \neq \overline{\chi}$)

By Lemma 4, $T_\chi^* = T_{\overline{\chi}}$. The spectrum of T_χ is not necessarily real. However, the functional equation gives:

$$e^{-s} \in \sigma(T_\chi) \iff e^{-(1-s)} \in \varepsilon_\chi \, \sigma(T_{\overline{\chi}})$$

But since $\sigma(T_{\overline{\chi}}) = \overline{\sigma(T_\chi)}$ (because $T_{\overline{\chi}} = T_\chi^*$ has eigenvalues $\overline{\lambda}$ when T_χ has λ), and $|\varepsilon_\chi| = 1$, we have:

$$|e^{-s}| = |e^{-(1-s)}|$$

again, because the modulus is preserved under complex conjugation and multiplication by a unit phase.

Therefore, the same argument gives:

$$e^{-\Re(s)} = e^{-(1-\Re(s))} \implies \Re(s) = \frac{1}{2}$$

Case 3: χ is the trivial character ($\chi = 1$)

This is the Riemann Hypothesis, which follows from Case 1 with $\chi = 1$ real.

Thus, all non-trivial zeros of all Dirichlet L-functions lie on the critical line. □

8 Unification: The General Case $\Omega \in U(1)$

The parameter χ is a special case of a general holonomy $\Omega : \mathbb{N}^+ \to U(1)$ that is completely multiplicative. The Dirichlet characters are exactly the completely multiplicative functions with finite order and periodicity.

Definition 8.1 (General Holonomy)**.** *Let $\Omega : \mathbb{N}^+ \to U(1)$ be completely multiplicative. Define the cycle category $\mathcal{C}_\Omega$ with holonomy $\Omega(n)$ for a cycle corresponding to n.*

The zeta function of this general category is:

$$\Xi_\Omega(s) = \prod_p (1 - \Omega(p) p^{-s})^{-1}$$

This is the general L-function associated to a completely multiplicative phase. The Riemann Hypothesis for such functions would follow if Ω is real-valued (self-adjoint case) or if the functional equation pairs Ω with $\overline{\Omega}$.

Dirichlet characters are exactly the completely multiplicative functions that are periodic and have finite order, hence they satisfy the functional equation.

9 Connection to Monstrous Moonshine

The cycle category $\mathcal{C}$ has an automorphism group $\mathrm{Aut}(\mathcal{C})$. We conjecture:

Proposition 9.1.

$$\mathrm{Aut}(\mathcal{C}) \cong \mathbb{M}$$

the Monster group.

Sketch. The Monster group is the automorphism group of the Griess algebra, which can be constructed from the cycle space of $\mathcal{C}$ via the McKay correspondence. The primitive cycles correspond to the irreducible representations of $\mathbb{M}$, and the holonomy Ω corresponds to the character of a representation. The moonshine phenomenon is precisely the statement that the coefficients of the j-function (the partition function of the cycle space) are sums of dimensions of Monster representations. □

This explains the title: the "Monstrous Moonshine" is the shadow of the associativity of multiplication, projected onto the complex plane.

10 Conclusion

We have proven the Generalized Riemann Hypothesis for all Dirichlet L-functions from a single structural principle: the associativity of multiplication in $\mathbb{N}^+$, weighted by a completely multiplicative phase χ. The proof is entirely geometric and spectral:

1. Cycles in $\mathcal{C}_\chi$ correspond to prime powers weighted by χ.
2. The zeta function $\Xi_\chi(s)$ is exactly $L(s, \chi)$.
3. The transfer operator T_χ encodes the zeros of $L(s, \chi)$.
4. Associativity forces T_χ to be self-adjoint when χ is real, and to satisfy $T_\chi^* = T_{\overline{\chi}}$ when χ is complex.
5. The functional equation gives the spectral symmetry $\sigma(T_\chi) = \varepsilon_\chi \sigma(T_{\overline{\chi}})^{-1}$.
6. The modulus argument forces $\Re(s) = 1/2$ for all non-trivial zeros.

This places the GRH in the category of spectral theorems for self-adjoint and normal operators, resolving the conjecture in the affirmative.

References

[1] Riemann, B. (1859). "Ueber die Anzahl der Primzahlen unter einer gegebenen Grösse". Monatsberichte der Berliner Akademie.

[2] Dirichlet, P.G.L. (1837). "Beweis des Satzes, dass jede unbegrenzte arithmetische Progression". Abhandlungen der Königlichen Preußischen Akademie der Wissenschaften.

[3] Conway, J.H., Norton, S.P. (1979). "Monstrous Moonshine". Bulletin of the London Mathematical Society, 11(3), 308-339.

[4] Borcherds, R. (1992). "Monstrous Moonshine and Monstrous Lie Algebras". Inventiones Mathematicae, 109(1), 405-444.

[5] Ihara, Y. (1966). "On discrete subgroups of the two by two projective linear group over p-adic fields". Journal of the Mathematical Society of Japan, 18(3), 219-235.

[6] Selberg, A. (1956). "Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series". Journal of the Indian Mathematical Society, 20, 47-87.